

Equation of state of a seven-dimensional hard-sphere fluid. Percus–Yevick theory and molecular-dynamics simulations

Miguel Robles^{a)} and Mariano López de Haro^{b)}

Centro de Investigación en Energía, UNAM, Temixco, Morelos 62580, Mexico

Andrés Santos^{c)}

Departamento de Física, Universidad de Extremadura, E-06071 Badajoz, Spain

(Received 14 August 2003; accepted 18 February 2004)

Following the work of Leutheusser [*Physica A* **127**, 667 (1984)], the solution to the Percus–Yevick equation for a seven-dimensional hard-sphere fluid is explicitly found. This allows the derivation of the equation of state for the fluid taking both the virial and the compressibility routes. An analysis of the virial coefficients and the determination of the radius of convergence of the virial series are carried out. Molecular-dynamics simulations of the same system are also performed and a comparison between the simulation results for the compressibility factor and theoretical expressions for the same quantity is presented. © 2004 American Institute of Physics.

[DOI: 10.1063/1.1701840]

I. INTRODUCTION

In liquid theory there has been a long lasting interest on the equilibrium properties of high-dimensional hard-sphere fluids, especially in the last few years.^{1–30} Such an interest has arisen from many different sources. To begin with, given the relative simplicity of the intermolecular interactions in these hard-core systems, they are amenable to both theoretical and computer simulation studies. In this sense and as it occurs in other problems in theoretical and mathematical physics, it is an asset that one can deal with hard spheres in arbitrary dimensionality and exploit some of the features that these systems have in common, for instance the fact that they all exhibit a first-order freezing transition. Furthermore, and as conjectured by Frisch and Percus,¹⁹ in the case of fluids high spatial dimensionality may have a parallel with limiting high density situations, so that by increasing the dimensionality one may obtain at least a rough idea of any thermodynamic phenomenology that extends to such dimensionality. An example of this expectation is the recent investigation of the demixing problem in mixtures of hard hyperspheres.²²

Computer simulation studies of hard-sphere fluids in dimensions greater than three are very scarce. To the best of our knowledge, only the four- and five-dimensional simple^{5,13} and multicomponent²⁵ fluids have been simulated. This is not surprising since the computational effort needed to obtain reliable results increases significantly with the dimensionality.

Exact information on the equation of state (EOS) usually comes from the virial coefficients B_n defined by³¹

$$Z \equiv \frac{p}{\rho k_B T} = 1 + \sum_{n=2}^{\infty} B_n \rho^{n-1} = 1 + \sum_{n=2}^{\infty} b_n \eta^{n-1}. \quad (1)$$

In this equation, p is the pressure, ρ is the number density, k_B

is the Boltzmann constant, T is the temperature, and Z is the compressibility factor. The second virial coefficient is $B_2 = 2^{d-1} v_d \sigma^d$, where d is the dimensionality, σ is the diameter of a sphere, and $v_d = (\pi/4)^{d/2} / \Gamma(1+d/2)$ is the volume of a d -dimensional sphere of unit diameter. In the second equality of Eq. (1) we have introduced the packing fraction $\eta \equiv \rho v_d \sigma^d$ and the reduced virial coefficients $b_n \equiv B_n / (v_d \sigma^d)^{n-1} = 2^{(d-1)(n-1)} B_n / B_2^{n-1}$. The radius of convergence of the virial series (1), $\rho_{\text{conv}} = \lim_{n \rightarrow \infty} |B_n / B_{n+1}|$, is the modulus of the singularity of $Z(\rho)$ closest to the origin in the complex ρ plane. If such a singularity were located on the positive real axis, then all the virial coefficients B_n would be positive for large n .

The exact expression for the third virial coefficient is^{2,9}

$$\frac{B_3}{B_2^2} = 2 \frac{B_{3/4}(d/2+1, 1/2)}{B(d/2+1/2, 1/2)}, \quad (2)$$

where $B(a, b) = \Gamma(a)\Gamma(b)/\Gamma(a+b)$ is the beta function and $B_x(a, b)$ is the incomplete beta function. Both B_2 and B_3 are positive definite for arbitrary d . The analytic evaluation of the fourth virial coefficient is much more involved. Luban and Baram^{2,3} derived exact expressions for two of the three terms contributing to B_4 and proposed a semiempirical formula for the remaining contribution. More recently, Clisby and McCoy^{28,30} showed that B_4 can be evaluated exactly for any even dimension d and gave the explicit results for $d = 4, 6, 8, 10$, and 12 . They also computed numerically^{29,30} the fifth and sixth virial coefficients through $d = 50$. The results show that, while B_5 remains positive, B_4 and B_6 become negative for $d \geq 8$ and $d \geq 6$, respectively. This suggests the possibility that, even in the three-dimensional case, there might be negative virial coefficients B_n for sufficiently large n .³² The fact that the virial coefficients are not positive definite and that they may have alternate signs is of importance in connection with the radius of convergence of the virial expansion (1), as mentioned before.

^{a)}Electronic mail: mrp@cie.unam.mx

^{b)}Electronic mail: malopez@servidor.unam.mx

^{c)}http://www.unex.es/fisteor/andres/. Electronic mail: andres@unex.es

A. The scenario for high dimensionality

The high-dimensionality limit of hard hyperspheres has been the subject of several studies.^{6,7,19,23} By means of asymptotic methods and heuristic arguments, Frisch and Percus¹⁹ were led to the following scenario in that limit:

- (i) The fourth virial coefficient is negative. Beyond that term, the virial expansion is an alternating series;
- (ii) the virial series is convergent for $\hat{\rho} < 1$, where $\hat{\rho} \equiv 2\eta^{1/d}$ is the scaled density per dimension. In terms of the packing fraction, the virial series converges for $\eta < \eta_{\text{conv}} = 2^{-d}$;
- (iii) in the density range $\hat{\rho} < 1$ the second virial term dominates over the remaining ones, so that

$$Z \approx 1 + B_2 \rho = 1 + \frac{\hat{\rho}^d}{2}. \quad (3)$$

- (iv) Even though the virial expansion does not converge for $\hat{\rho} > 1$ (oscillatory divergence), the truncated series (3) remains valid within the interval $1 < \hat{\rho} < (1 - \epsilon)\hat{\rho}_0$, where $\epsilon = \mathcal{O}(d^{-1})$ and

$$\hat{\rho}_0 = (1.148d^{-1/6}e^{-1.473d^{1/3}})^{-1/d} \sqrt{e/2} \sim \sqrt{e/2} \approx 1.17. \quad (4)$$

- (v) At the density $\hat{\rho} = \hat{\rho}_0$ an infinite compressibility spinodal appears, thus indicating a first-order transition to the high-dimensional solid.

In an independent paper, Parisi and Slanina²³ reached similar conclusions from a toy model based on simplified HNC equations. They obtained that, while in the limit $d \rightarrow \infty$ the EOS for $\hat{\rho} < 1$ is given by Eq. (3), in the interval $1 < \hat{\rho} < \hat{\rho}_0 = \sqrt{e/2}$, one has

$$Z(\hat{\rho}) = 1 + \hat{\rho}^d \left[\frac{1}{2} + \Delta(\hat{\rho}) \right], \quad (5)$$

where

$$\Delta(\hat{\rho}) = \left[\frac{e\kappa(\hat{\rho})}{2\hat{\rho}^2} \right]^d, \quad (6)$$

$\kappa(\hat{\rho})$ being the solution to³³

$$\ln \frac{2\hat{\rho}^2}{e} = \ln(1 + \sqrt{1 - \kappa^2}) - \sqrt{1 - \kappa^2}. \quad (7)$$

Note that, since $\ln \kappa < \ln(1 + \sqrt{1 - \kappa^2}) - \sqrt{1 - \kappa^2}$ for $0 < \kappa < 1$, one has $\lim_{d \rightarrow \infty} \Delta(\hat{\rho}) = 0$ for $\hat{\rho} < \hat{\rho}_0$. Although, strictly speaking, Eq. (7) cannot be extended to $\hat{\rho} > \hat{\rho}_0$, Eq. (6) suggests that $\lim_{d \rightarrow \infty} \Delta(\hat{\rho}) = \infty$ in that density domain, in agreement with the phase transition noted by Frisch and Percus.³⁴

B. Approximate equations of state

As in two and three dimensions, one can make use of approximate schemes to represent the EOS of hard hyperspheres. Several proposals have been made in the literature for the EOS based on the knowledge of the first few virial coefficients.^{9–12,15} For illustration, we review here a few of them making use of the first three virial coefficients. The extension to higher virial coefficients is straightforward.

1. Truncated virial series

The first obvious choice is the truncated virial expansion

$$Z_{[2,0]}(\eta) = 1 + b_2 \eta + b_3 \eta^2. \quad (8)$$

As discussed above, this simple approximation becomes more and more accurate in the stable fluid domain as the dimensionality increases. In a way analogous to Eq. (8) it is possible to define a truncated expansion $Z_{[n,0]}$ from the knowledge of the first $n+1$ virial coefficients.

2. Padé approximants

One can also construct Padé approximants of the form $Z_{[n,m]}$ from the first $n+m+1$ virial coefficients. For instance,

$$Z_{[1,1]}(\eta) = \frac{b_2 + (b_2^2 - b_3)\eta}{b_2 - b_3\eta}, \quad (9)$$

$$Z_{[0,2]}(\eta) = [1 - b_2\eta + (b_2^2 - b_3)\eta^2]^{-1}. \quad (10)$$

3. Colot–Baus approximation

Colot and Baus^{8,9} proposed (truncated) *rescaled* virial expansions, where the series expansion of $(1 - \eta)^d Z(\eta)$, rather than that of $Z(\eta)$, is truncated. Let us denote by $Z_{[n,0]}^{\text{BC}}(\eta)$ the truncated rescaled virial expansion that makes use of the first $n+1$ virial coefficients. For example,

$$Z_{[2,0]}^{\text{BC}}(\eta) = \frac{1 + (b_2 - d)\eta + [b_3 - b_2d + d(d-1)/2]\eta^2}{(1 - \eta)^d}. \quad (11)$$

The pole of order d at the (unphysical) packing fraction $\eta = 1$ is suggested by the scaled particle theory.

4. Maeso–Solana–Amorós–Villar (MSAV) approximation

Maeso *et al.*¹⁵ combined the advantages of Padé approximants and rescaled expansions by proposing Padé approximants for $(1 - \eta)^d Z(\eta)$. A rescaled Padé approximant constructed from the first $n+m+1$ virial coefficients will be denoted here as $Z_{[n,m]}^{\text{MSAV}}(\eta)$. Thus,

$$Z_{[1,1]}^{\text{MSAV}}(\eta) = \frac{1}{(1 - \eta)^d} \times \frac{b_2 - d + [d(d+1)/2 + b_2(b_2 - d) - b_3]\eta}{b_2 - d - [b_3 - b_2d + d(d-1)/2]\eta}. \quad (12)$$

By construction, $Z_{[n,0]}^{\text{MSAV}}(\eta) = Z_{[n,0]}^{\text{BC}}(\eta)$.

5. Song–Mason–Stratt approximation

Using simple arguments, Song *et al.*^{10,12} proposed the following generalization to d dimensions of the celebrated Carnahan–Starling (CS) EOS for three-dimensional hard spheres:³⁵

$$Z_{\text{SMS}}(\eta) = 1 + b_2 \eta \frac{1 + (b_3/b_2 - d)\eta}{(1 - \eta)^d}. \quad (13)$$

6. Luban–Michels approximation

On a different vein, Luban and Michels¹³ wrote the compressibility factor as

$$Z_{LM}(\eta) = 1 + b_2 \eta \times \frac{1 + [b_3/b_2 - \zeta(\eta)b_4/b_3]\eta}{1 - \zeta(\eta)(b_4/b_3)\eta + [\zeta(\eta) - 1](b_4/b_2)\eta^2}. \quad (14)$$

The knowledge of the function $\zeta(\eta)$ is equivalent to that of $Z(\eta)$. However, $\zeta(\eta)$ focuses on the high density behavior of the EOS, since Eq. (14) is consistent with the exact first four virial coefficients, regardless of the choice of $\zeta(\eta)$. The approximation $\zeta(\eta)=1$ is equivalent to assuming a Padé approximant $Z_{[2,1]}(\eta)$. Instead, Luban and Michels observed that the computer simulation data for $d=2-5$ favor a linear approximation $\zeta(\eta)=a+b\eta$, with coefficients obtained by a least-square fit to the simulation results for each dimensionality.

7. Percus–Yevick theory

It is noteworthy that the Percus–Yevick (PY) integral equation can be solved analytically in odd dimensions, as first pointed out by Freasier and Isbister¹ and, independently, Leutheusser.⁴ The latter concluded that, in general, the problem reduces to an algebraic equation of degree $d-3$. Following his procedure, however, we find that for $d \geq 9$ this is not so (see the Appendix) and our calculations suggest that such degree should rather be $2^{(d-3)/2}$ for $d \geq 3$. In any case, in five dimensions one has to deal with a quadratic equation^{1,4} and explicit expressions for the virial and compressibility routes to the EOS can be obtained.²⁰ A simple analysis of the solution for $d=5$, that as far as we know has not been carried out before, shows that the virial route incorrectly gives a negative value for B_6 : $B_6^{\text{PY-v}}/B_2^5 = -2999/16^5 \approx -0.00286$. The compressibility route yields $B_6^{\text{PY-c}}/B_2^5 = 12\,233/(8 \times 16^5) \approx 0.00146$, while the correct value is $B_6/B_2^5 \approx 0.00094$.^{29,30} Both routes consistently predict that B_8 is negative, with subsequent coefficients alternating in sign. On the other hand, the virial route gives values for the magnitude of B_n ($n \geq 8$) increasingly larger than the compressibility route: $B_n^{\text{PY-v}}/B_n^{\text{PY-c}} \approx 0.66 + 0.77n$. The alternating character of the virial series predicted by the PY equation for $d=5$ is due to a branch singularity located on the *negative* real axis at $\eta_{\text{branch}} = -(9 - 5\sqrt{3})/6 \approx -0.0566243$. The radius of convergence $\eta_{\text{conv}}^{\text{PY}} \approx 0.0566243$ of the PY solution for $d=5$ is larger than the value $2^{-5} = 0.03125$ extrapolated from the radius $\lim_{d \rightarrow \infty} \eta_{\text{conv}} = 2^{-d}$, but is close to the estimate $\eta_{\text{conv}} \approx 0.052$ made by Clisby and McCoy on the basis of Monte Carlo evaluation of sets of Ree–Hoover diagrams.^{29,30} All these estimates are sensibly smaller than the packing fraction $\eta_f = 0.19$ at which freezing occurs for $d=5$.^{5,13,26}

8. Generalized Carnahan–Starling approximation

As is well known, the CS EOS for three-dimensional hard spheres can be interpreted as a weighted average between the PY virial and compressibility routes:

$$Z_{CS}(\eta) = \alpha Z_{\text{PY-c}}(\eta) + (1 - \alpha) Z_{\text{PY-v}}(\eta), \quad (15)$$

where $\alpha=2/3$. Given that the PY equation can be solved for odd dimensions, it is then natural to speculate about whether or not the prescription (15), with an adequate choice of the mixing parameter α , keeps being reliable for $d > 3$, even though the internal inconsistency between both routes seems to increase dramatically with the dimensionality.¹ In the five-dimensional case, one of us²¹ showed that the choice $\alpha=3/5$ leads to values of Z_{CS} in excellent agreement with computer simulations.⁵ This suggested that the choice $\alpha=(d+1)/2d$ might provide a good description for $d \geq 3$. Note that, while Eqs. (13) and (15) coincide at $d=3$, they differ for $d > 3$, so they generalize the original CS EOS along different directions. An alternative generalization of the CS EOS was made by Gonzalez *et al.*¹⁶ They proposed a simple ansatz for the direct correlation function $c(r)$, which reduced exactly to the PY theory for $d=1$ and $d=3$ and gave results very close to the PY theory for other dimensions. Their generalized CS EOS consisted of a weighted average between the virial and compressibility routes obtained from their theory with a mixing parameter $\alpha=2(2d-1)/5d$.

C. Aim of the paper

The aim of this paper is threefold. First, we present the explicit solution to the PY equation in the case of a seven-dimensional hard-sphere fluid following the procedure introduced by Leutheusser.⁴ This allows us to derive the EOS of the fluid both through the virial and the compressibility routes, as well as to analyze the behavior of the virial coefficients stemming out of them. As we will see, the singularity closest to the origin is again a branch point on the negative real axis, so the radius of convergence of the PY virial series is $\eta_{\text{conv}} \approx 0.0100625$. We conjecture that this value might be close to the (unknown) exact radius. Moreover, a Carnahan–Starling-type equation of state of the form (15) with $\alpha=5/6$ is proposed. Secondly, we provide molecular-dynamics results for the compressibility factor. To the best of our knowledge, this is the first time that simulation results are presented for hard hyperspheres in seven dimensions. The twenty densities considered range from the dilute regime ($\rho\sigma^7=0.1$ or $\eta=0.0037$) to our estimated freezing point ($\rho\sigma^7 \approx 0.95$ or $\eta \approx 0.072$). Finally, we perform a comparison between different proposals for the EOS of a seven-dimensional hard-sphere fluid with the simulation data. We observe that the proposals (11) and (13) (which do not have any empirical parameter), (14) (which contains two fitting parameters), and (15) (with one fitting parameter) reproduce fairly well the simulation data.

The paper is organized as follows. In Sec. II we provide the solution of the PY equation for a seven-dimensional hard-sphere fluid as well as the analysis of the virial coefficients arising from the derivation of the EOS using the virial and the compressibility routes. This is followed in Sec. III by a description of the molecular-dynamics simulation that was carried out to obtain the compressibility factor of the fluid. The results of the simulation are then used to assess the merits of various proposals that have been made in the literature for the EOS. The paper is closed in Sec. IV with

TABLE I. Values of B_n/B_2^{n-1} for $n=3-8$, according to the virial route of the PY approximation, the compressibility route of the PY approximation, the CS-type approximation (15) with $\alpha=5/6$, and the known exact results (Refs. 29 and 30).

n	$B_n^{\text{PY-v}}/B_2^{n-1}$	$B_n^{\text{PY-c}}/B_2^{n-1}$	$B_n^{\text{CS}}/B_2^{n-1}$	$B_n^{\text{ex}}/B_2^{n-1}$
3	0.282 226 562 5	0.282 226 562 5	0.282 226 562 5	0.282 226 562 5
4	$-7.499\ 694\ 824 \times 10^{-3}$	$2.155\ 081\ 431 \times 10^{-2}$	$1.670\ 906\ 279 \times 10^{-2}$	$9.873(4) \times 10^{-3}$
5	$1.235\ 022\ 893 \times 10^{-2}$	$5.116\ 807\ 918 \times 10^{-3}$	$6.322\ 378\ 086 \times 10^{-3}$	$7.071(7) \times 10^{-3}$
6	$-8.177\ 005\ 666 \times 10^{-3}$	$-1.865\ 328\ 120 \times 10^{-3}$	$-2.917\ 274\ 378 \times 10^{-3}$	$-3.52(2) \times 10^{-3}$
7	$6.553\ 131\ 160 \times 10^{-3}$	$1.384\ 246\ 670 \times 10^{-3}$	$2.245\ 727\ 418 \times 10^{-3}$	
8	$-5.762\ 797\ 816 \times 10^{-3}$	$-1.078\ 783\ 146 \times 10^{-3}$	$-1.859\ 452\ 258 \times 10^{-3}$	

further discussion of the results and some concluding remarks.

II. SOLUTION OF THE PERCUS–YEVICK EQUATION FOR A SEVEN-DIMENSIONAL HARD-SPHERE FLUID

As mentioned in Sec. I, the solution to the PY equation for hard hyperspheres with $d=\text{odd}$ reduces to an algebraic equation of degree $2^{(d-3)/2}$. The case $d=5$, which yields a quadratic equation, has been analyzed by several authors.^{1,4,14,20,21} The highest dimensionality for which the algebraic problem certainly lends itself to an analytic solution is $d=7$. A sketch of the general solution and some details of the particular cases $d=7$ and $d=9$ are provided in the Appendix. It is shown there that the solution of the PY equation for seven-dimensional hard hyperspheres is given by the physical solution to the quartic equation (A19). In the Appendix it is also shown that for $d=9$ the resulting algebraic equation is of eighth degree.

A study of the solutions of Eq. (A19) shows that in the interval $0.446\ 469 \leq \eta < 1$ the four roots are real. On the other hand, for $0 \leq \eta \leq 0.446\ 469$ two of the roots become complex conjugates and only the other two roots remain real, the physical one being finite in the limit $\eta \rightarrow 0$. The explicit solution to Eq. (A19) involves the term $[P_4(\eta)P_6(\eta)]^{1/2}$, where $P_4(\eta) = 1 + 94\eta + 202\eta^2 + \frac{1360}{3}\eta^3 + 50\eta^4$ and $P_6(\eta) = 1 + 99\eta - \frac{307}{8}\eta^2 - \frac{339}{4}\eta^3 - \frac{2762}{3}\eta^4 + \frac{695}{2}\eta^5 + \frac{5575}{108}\eta^6$. As a consequence, the solution possesses branch points at the zeroes of $P_4(\eta)$ and $P_6(\eta)$. The zero of $P_4(\eta)$ closest to the origin is $\eta'_{\text{branch}} \approx -0.010\ 886\ 8$, while that of $P_6(\eta)$ is $\eta_{\text{branch}} \approx -0.010\ 062\ 5$. Therefore, the radius of convergence of the virial series for a seven-dimensional hard-sphere fluid described by the PY approximation is $\eta_{\text{conv}}^{\text{PY}} = |\eta_{\text{branch}}| \approx 0.010\ 062\ 5$.

Table I gives the first few values of the PY virial coefficients obtained from the virial and the compressibility routes. As far as we know, the exact values B_n^{ex} of the virial coefficients of seven-dimensional hard spheres are known up to $n=6$ only.^{29,30} They are listed in Table I as well, which also gives the CS-type values $B_n^{\text{CS}}/B_2^{n-1}$, where $B_n^{\text{CS}} = \alpha B_n^{\text{PY-c}} + (1-\alpha)B_n^{\text{PY-v}}$ with the simple choice $\alpha=5/6$. Note that the choice $\alpha \approx 0.6$ would make $B_4^{\text{CS}} \approx B_4^{\text{ex}}$, whereas the choice $\alpha \approx 0.7$ would make $B_5^{\text{CS}} \approx B_5^{\text{ex}}$ and $B_6^{\text{CS}} \approx B_6^{\text{ex}}$. However, comparison with molecular-dynamics simulations (see Sec. III) favors $\alpha \approx 0.8$.

From Table I we observe that the virial route of the PY approximation incorrectly yields a negative value for the

fourth virial coefficient (which actually becomes negative for $d \geq 8$,²⁸⁻³⁰) while the compressibility route predicts the correct sign.³⁶ We have computed $B_n^{\text{PY-v}}$ and $B_n^{\text{PY-c}}$ for values of n much larger than those displayed in Table I. The results indicate that $\text{sgn}(B_n^{\text{PY-v}}) = (-1)^{n+1}$ for $5 \leq n \leq 97$ but $\text{sgn}(B_n^{\text{PY-v}}) = (-1)^n$ for $n \geq 98$; analogously, $\text{sgn}(B_n^{\text{PY-c}}) = (-1)^{n+1}$ for $5 \leq n \leq 80$ but $\text{sgn}(B_n^{\text{PY-c}}) = (-1)^n$ for $n \geq 81$. Therefore, both routes synchronize their signs for $5 \leq n \leq 80$ and again for $n \geq 98$. This peculiar behavior of the alternating character of the virial series seems to be a consequence of the proximity between the two branch point singularities closest to the origin, $\eta_{\text{branch}} \approx -0.010\ 062\ 5$ and $\eta'_{\text{branch}} \approx -0.010\ 886\ 8$, both located on the negative real axis. To confirm this interpretation, we plot in Fig. 1 the ratios $|b_n^{\text{PY-v}}/b_{n+1}^{\text{PY-v}}|$, $|b_n^{\text{PY-c}}/b_{n+1}^{\text{PY-c}}|$, and $|b_n^{\text{CS}}/b_{n+1}^{\text{CS}}|$. Recall that the radius of convergence of the virial series is $\eta_{\text{conv}} = \lim_{n \rightarrow \infty} |b_n/b_{n+1}|$. Figure 1 shows that for $n \leq 50$ the ratio $|b_n/b_{n+1}|$ seems to converge from above to the apparent radius of convergence $\eta'_{\text{conv}} = -\eta'_{\text{branch}} \approx 0.010\ 886\ 8$. However, the true radius $\eta_{\text{conv}}^{\text{PY}} = -\eta_{\text{branch}} \approx 0.010\ 062\ 5$ is reached from below for $n \geq 100$.

As mentioned in Sec. I, the radius of convergence predicted by the PY approximation in the five-dimensional case is $\eta_{\text{conv}}^{\text{PY}} \approx 0.056\ 624\ 3$. When going to the next odd dimensionality, the radius of convergence has shrunk to $\eta_{\text{conv}}^{\text{PY}} \approx 0.010\ 062\ 5$. In terms of the scaled density per dimension introduced by Frisch and Percus,¹⁹ the radius of convergence is $\hat{\rho}_{\text{conv}}^{\text{PY}} \approx 1.126$ for $d=5$ and $\hat{\rho}_{\text{conv}}^{\text{PY}} \approx 1.037$ for $d=7$. Therefore, it nicely tends to converge to the expected value $\hat{\rho}_{\text{conv}}$




FIG. 1. Plot of the ratios $|b_n^{\text{PY-c}}/b_{n+1}^{\text{PY-c}}|$ (diamonds), $|b_n^{\text{PY-v}}/b_{n+1}^{\text{PY-v}}|$ (triangles), and $|b_n^{\text{CS}}/b_{n+1}^{\text{CS}}|$ (circles). The horizontal lines correspond to the apparent radius of convergence $\eta'_{\text{conv}} \approx 0.010\ 886\ 8$ and to the true radius of convergence $\eta_{\text{conv}}^{\text{PY}} \approx 0.010\ 062\ 5$.




FIG. 2. Plot of the virial coefficients B_n^{PY-c}/B_2^{n-1} (diamonds), B_n^{PY-v}/B_2^{n-1} (triangles), and B_n^{CS}/B_2^{n-1} (circles).

$= 1$ as $d \rightarrow \infty$. In addition, the PY value of $\eta_{conv} = 2^{1-d} B_2 \lim_{n \rightarrow \infty} |B_n/B_{n+1}|$ for $d=7$ is consistent with estimates obtained from Table XVIII of Ref. 29. By assuming that the Ree–Hoover ring diagrams dominate for high d ,²⁹ one has $\eta_{conv} < 2^{-6} B_2 |B_9/B_{10}| \approx 2^{-6} 0.0132/0.0143 \approx 0.014$ for $d=7$, which agrees with the PY value $\eta_{conv}^{PY} \approx 0.010\,062\,5$. Clisby and McCoy’s estimate²⁹ $\eta_{conv} \approx 0.052$ for $d=5$ is also close to the PY value $\eta_{conv}^{PY} \approx 0.056\,624\,3$. All of this leads us to conjecture that the PY solution gives a fair estimate of the radius of convergence of the true virial series for high dimensionalities. Pushing this conjecture even further, we can expect the true radius of convergence to be due to a singularity (pole or branch point) located on the negative real axis, so that the virial coefficients alternate in sign beyond a certain order. Figure 2 shows the virial coefficients B_n^{PY-v}/B_2^{n-1} , B_n^{PY-c}/B_2^{n-1} , and B_n^{CS}/B_2^{n-1} in the seven-dimensional case. In the spirit of the above conjecture, one may speculate that the exact values of B_n/B_2^{n-1} lie in between B_n^{PY-v}/B_2^{n-1} and B_n^{PY-c}/B_2^{n-1} , perhaps not far from the interpolated values B_n^{CS}/B_2^{n-1} . The reduced virial coefficients B_n/B_2^{n-1} start decreasing in magnitude, reach a minimum around $n=10$, and then grow with n . The fact that the PY solution in the three-dimensional case does not possess a branch point singularity, so that all the virial coefficients remain positive, casts some doubts as to whether the true virial series fails to converge for densities close to the freezing density $\eta_f \approx 0.494$. In any case, the true radius of convergence for $d=3$ cannot be larger than the crystalline close-packing value $\eta_{cp} = \pi\sqrt{2}/6 \approx 0.7405$, while the PY solution has $\eta_{conv}^{PY} = 1$.

III. MOLECULAR-DYNAMICS SIMULATIONS

A. Method

The numerical simulation was implemented by using the same algorithm as described in Ref. 25, which is also based on the work of Michels and Trappeniers⁵ and Luban and Michels¹³ for four- and five-dimensional hyperspheres. We are not aware of any previous computer simulation of hard hyperspheres of a dimension higher than $d=5$. We have chosen the molecular-dynamics method instead of the Monte Carlo method because that gives us the possibility of testing our code by applying it to $d=4$ and $d=5$ and comparing with the results of Refs. 5 and 13.

For our simulations, in order to keep the computing time within reasonable limits and at the same time being able to examine a wide density range, the initial configuration is chosen to be the one obtained by placing $N=64$ hyperspheres in a unitary cell of a d -type lattice. The simulation cell is a hypercube of side L and volume $V=L^d=N/\rho$ and the minimum image convention and periodic boundary conditions in all directions have been applied, in the same way as in the three-dimensional (3D) case.³⁷

During the simulations only binary collisions are taken into account, while collisions between three or four particles are ignored. The collision time for every pair of particles is calculated and the smallest value is obtained. All the particles are moved during this time at constant velocity. The pair of particles that suffers a collision is treated according to impulsive dynamics and the velocities are changed; in this step the hard collisional virial is calculated. This allows one to evaluate the excess compressibility factor as

$$Z - 1 = - \frac{1}{N \langle v^2 \rangle \Delta t} \sum_{ij} \mathbf{r}_{ij} \cdot \Delta \mathbf{v}_i, \quad (16)$$

where $\langle v^2 \rangle$ is the mean-square velocity, Δt is the simulated time, \mathbf{r}_{ij} is the relative position vector between colliding particles i and j , and $\Delta \mathbf{v}_i$ the change in velocity of the particle i on collision.

The equation of state is achieved by changing the diameter σ of the particles, in such a way that the reduced density $\rho^* = \rho \sigma^d$ changes, and letting the system to relax up to an equilibrium pressure. The errors associated with our calculation were computed following standard methods for errors in equilibrium averages.³⁷

As mentioned above, before running the program for $d=7$, it has been previously validated for $d=4$ and $d=5$, reproducing the excess compressibility factor obtained by Michels, Trappeniers, and Luban.^{5,13}

B. Results

We have computed the compressibility factor for densities $0.1 \leq \rho^* \leq 1.90$ with a step $\Delta \rho^* = 0.1$, as well as for $\rho^* = 1.95$. The simulation data obtained by our molecular-dynamics simulations are listed in Table II. At the largest density $\rho^* = 1.95$ ($\eta = 0.0720$) the compressibility factor presents a dramatic drop. We interpret this as an indication of the freezing transition. Consequently, the density at which the seven-dimensional fluid of hard spheres freezes can be estimated as $\rho_f^* \approx 1.95$ or, equivalently, $\eta_f \approx 0.072$. From Fig. 5 of Ref. 26 one can observe that $\ln \eta_f(d)$ is almost a linear function of the dimensionality d , with a slight negative curvature. According to this, knowing the freezing densities $\eta_f(d)$ and $\eta_f(d+2)$, one can estimate the freezing density $\eta_f(d+4)$ as $\eta_f(d+4) \approx \eta_f^2(d+2)/\eta_f(d)$. Given that $\eta_f(3) \approx 0.494$ and $\eta_f(5) \approx 0.19$, one has $\eta_f(7) \approx 0.19^2/0.494 \approx 0.073$, in close agreement with our estimate. An independent estimate based on a conjecture by Colot and Baus⁸ confirms again this value. These authors suggested that the ratio of length scales $[\eta_f(d)/\eta_{cp}(d)]^{1/d}$ is practically independent of d , so that $\eta_f(d+2) \approx \eta_{cp}(d+2)[\eta_f(d)/\eta_{cp}(d)]^{(d+2)/d}$. The general expression for the close-packing fraction $\eta_{cp}(d)$ is not known, but for $d < 25$ the values are

TABLE II. Compressibility factor as a function of η from the simulation data and for different approximations. The numbers in parentheses indicate the statistical error in the last significant digit.

η	Z_{simul}	Z_{CS}	$Z_{\text{PY-v}}$	$Z_{\text{PY-c}}$	$Z_{[4,0]}$	$Z_{[2,2]}$	$Z_{[3,2]}$	$Z_{[2,0]}^{\text{BC}}$	$Z_{[2,2]}^{\text{MSAV}}$	Z_{SMS}	Z_{LM}
0.0037	1.253 66(2)	1.252 23	1.251 92	1.252 29	1.252 14	1.252 14	1.252 14	1.252 33	1.252 14	1.252 33	1.252 17
0.0074	1.5337(1)	1.537 51	1.535 16	1.537 97	1.536 87	1.536 87	1.536 81	1.538 29	1.536 85	1.538 27	1.537 25
0.0111	1.8646(3)	1.857 67	1.849 97	1.859 21	1.855 77	1.855 76	1.855 34	1.860 08	1.855 62	1.860 03	1.857 45
0.0148	2.2103(3)	2.214 82	2.196 91	2.218 40	2.210 93	2.210 94	2.209 30	2.220 06	2.210 33	2.219 93	2.215 65
0.0185	2.6174(2)	2.611 24	2.576 74	2.618 14	2.604 99	2.605 13	2.600 47	2.620 71	2.603 28	2.620 47	2.615 24
0.0221	3.0650(4)	3.049 46	2.990 39	3.061 28	3.041 11	3.041 70	3.030 80	3.064 67	3.037 16	3.064 23	3.0600
0.0258	3.5449(5)	3.532 19	3.438 90	3.550 85	3.522 98	3.524 81	3.502 43	3.554 69	3.515 02	3.553 99	3.553 94
0.0295	4.0989(7)	4.062 34	3.923 42	4.090 12	4.054 80	4.059 48	4.017 69	4.093 72	4.040 37	4.092 64	4.101 09
0.0332	4.7013(5)	4.643 02	4.445 19	4.682 58	4.641 33	4.651 83	4.579 11	4.684 83	4.617 13	4.683 26	4.705 19
0.0369	5.389(1)	5.277 57	5.005 55	5.331 98	5.287 85	5.309 24	5.189 44	5.331 30	5.249 71	5.329 10	5.369 29
0.0406	6.051(1)	5.969 55	5.605 92	6.042 28	6.000 15	6.040 73	5.851 64	6.036 59	5.943 07	6.033 58	6.095 11
0.0443	6.8179(6)	6.722 76	6.247 77	6.817 75	6.784 56	6.857 31	6.568 96	6.804 33	6.702 73	6.800 33	6.882 28
0.0480	7.6325(1)	7.541 21	6.932 69	7.662 92	7.647 94	7.772 55	7.344 90	7.638 37	7.534 89	7.633 17	7.727 27
0.0517	8.5133(2)	8.429 22	7.662 33	8.582 59	8.597 69	8.803 33	8.183 28	8.542 78	8.446 45	8.536 13	8.622 20
0.0554	9.4294(3)	9.391 34	8.438 41	9.581 92	9.641 72	9.970 83	9.088 29	9.521 86	9.445 19	9.513 48	9.553 55
0.0591	10.492(1)	10.4324	9.262 75	10.6664	10.7885	11.3020	10.0645	10.5801	10.5398	10.5697	10.5009
0.0628	11.570(3)	11.5577	10.1372	11.8417	12.0469	12.8317	11.1168	11.7224	11.7400	11.7096	11.4365
0.0664	12.694(1)	12.7725	11.0638	13.1142	13.4266	14.6056	12.2507	12.9537	13.0569	12.9381	12.3252
0.0701	13.907(3)	14.0828	12.0446	14.4904	14.9374	16.6847	13.4722	14.2794	14.5029	14.2607	13.1265
0.0720	9.039 44(6)	14.7756	12.5559	15.2196	15.7454	17.8637	14.1178	14.9794	15.2786	14.9589	13.4810

not far from Blichfeldt's upper estimate²⁶ $\eta_{\text{cp}}(d) \leq 2^{-d/2}(d+2)/2$. Using $\eta_f(5) \approx 0.19$, we get $\eta_f(7) \approx 0.076$, which is again consistent with our estimate.

Table II also gives some theoretical values: the PY predictions $Z_{\text{PY-v}}$ and $Z_{\text{PY-c}}$, the CS-type interpolation (15) with $\alpha=5/6$, the truncated virial expansion $Z_{[4,0]}$, the Padé approximants $Z_{[2,2]}$ and $Z_{[3,2]}$ [the three latter being obvious extensions of the approximations (8)–(10)], the rescaled virial expansion $Z_{[2,0]}^{\text{BC}}$ defined by Eq. (11), the rescaled Padé approximant $Z_{[2,2]}^{\text{MSAV}}$ defined by a natural extension of Eq. (12), the SMS approximation (13), and the LM proposal (14).

Although the knowledge of the sixth virial coefficient B_6 would allow one to consider the truncated series $Z_{[5,0]}$, it is not included in Table II because it turns out to be clearly inferior to $Z_{[4,0]}$. This is a consequence of the fact that $B_6 < 0$, so that $Z_{[5,0]} < Z_{[4,0]}$, while for small and moderate densities $Z_{[4,0]} < Z_{\text{simul}}$. This is a strong indication that the unknown seventh virial coefficient B_7 must be positive. Among the different Padé approximants that can be constructed from the knowledge of the first six virial coefficients, the best agreement with the simulation data is presented by $Z_{[2,2]}$ for $\rho^* \leq 1.4$ ($\eta \leq 0.0517$) and by $Z_{[3,2]}$ for $\rho^* \geq 1.4$ ($\eta \geq 0.0517$). It is interesting to note that both Padé approximants have poles on the negative real axis (at $\eta \approx -0.079$ in the case of $Z_{[2,2]}$ and at $\eta \approx -0.025$ in the case of $Z_{[3,2]}$), so that the extrapolated virial coefficients have alternating signs. Paradoxically, while the rescaled expansion $Z_{[2,0]}^{\text{BC}}$ incorporates the first three virial coefficients only, it exhibits a better agreement with simulation than those rescaled expansions that can be constructed with the first four, five, or six virial coefficients, so the latter are not included in Table II. Analogously, the best performance among the rescaled Padé approximants corresponds to $Z_{[2,2]}^{\text{MSAV}}$. Interestingly, the SMS proposal [cf. Eq. (13)] and the approximation $Z_{[2,0]}^{\text{BC}}$ yield practically equivalent results. The difference between both EOS is

$$Z_{[2,0]}^{\text{BC}}(\eta) - Z_{\text{SMS}}(\eta) = \frac{\eta^3}{(1-\eta)^7} (35 - 35\eta + 21\eta^2 - 7\eta^3 + \eta^4). \quad (17)$$

This corresponds to a relative difference smaller than 0.14% for the density range considered in the simulations.

Two of the theoretical EOS included in Table II, namely Z_{CS} and Z_{LM} , have an empirical character. The proposal (15) is based on the observation that the two PY routes tend to bracket the simulation data, as happens in the three-dimensional³⁵ and five-dimensional²¹ cases. We have found that the value $\alpha=5/6$ of the parameter is the simplest rational number that makes Z_{CS} reproduce fairly well the simulation values. In the case of the Luban–Michels EOS (14) one fits $\zeta(\eta)$ to a linear function. Figure 3 shows the simulation values of $\zeta(\eta)$. As in the five-dimensional case,¹³ $\zeta(\eta)$ is an increasing function of η , while it is a decreasing function for $d=2-4$. A linear fit in the interval $0.5 \leq \rho^* \leq 1.9$ ($0.0185 \leq \eta \leq 0.0701$) yields




FIG. 3. Plot of the simulation values of the function $\zeta(\eta)$ defined by Eq. (14). The dashed line is the linear fit (18).




FIG. 4. Compressibility factor as a function of the packing fraction. The circles represent simulation data, the dashed line represents $Z_{PY,c}$, the dash-dotted line represents $Z_{PY,v}$, and the solid line represents the CS-type interpolation (15) with $\alpha=5/6$.

$$\zeta(\eta) = -5.81 + 88.2\eta. \quad (18)$$

The column labeled Z_{LM} in Table II has been evaluated using the fit (18). On the other hand, our simulation data in Fig. 3 seem to indicate a negative curvature of $\zeta(\eta)$.

Table II shows that up to $\rho^*=0.8$ ($\eta=0.0295$) all the different theoretical results tabulated, including the simple truncated virial expansion $Z_{[4,0]}$, behave relatively well. For larger densities, $Z_{[4,0]}$ tends to overestimate the simulation data, while the Padé approximants $Z_{[2,2]}$ and $Z_{[3,2]}$ tend to underestimate them. The best global agreement is presented by Z_{CS} , Z_{LM} , $Z_{[2,0]}^{BC}$, and Z_{SMS} . This is especially noteworthy in the case of the two latter approximations, since they do not contain fitting parameters and, moreover, only the knowledge of the first three virial coefficients is exploited. This contrasts with Z_{LM} , which includes the fourth virial coefficient and contains two fitting parameters. On the other hand, Z_{CS} belongs in a different class of approximations. Given the involved algebraic structure of the PY solution, Z_{CS} does not intend to represent a practical recipe to the EOS of a seven-dimensional hard-sphere fluid. Instead, its role is to highlight the fact that the two PY routes keep bracketing the simulation data, so that an interpolation between them with a density-independent parameter α is rather accurate, as graphically illustrated in Fig. 4. This gives some confidence on the expectation that some of the analytical properties of the PY solution (e.g., alternating character of the virial series, branch points located on the negative real axis, ...) may shed light on the true behavior of the exact series.

IV. DISCUSSION

The results of the previous sections deserve further discussion. To begin with, to our knowledge this is the first time that a molecular dynamics simulation has been carried out on a seven-dimensional hard-sphere fluid. The simulation strategy that we adopted implied a compromise between computer process time and density range to be explored and the outcome is rather encouraging. The availability of simulation data for the EOS of the fluid allowed us to locate the freezing transition and also to assess the merits and limitations of

various proposals that have been made in the literature for the compressibility factor of hard hyperspheres. From this analysis it is clear that even simple approximations such as $Z_{[2,0]}^{BC}$ and Z_{SMS} do a reasonably good job and that, as it occurs in other dimensionalities, the virial and compressibility routes to the EOS in the PY approximation keep bracketing the simulation data, so that a Carnahan–Starling-type recipe of the form of Eq. (15) turns out to be rather accurate. However, the parameter α seems not to follow a simple relation as the ones suggested by González *et al.*^{16,17} or Santos.²¹

We also presented the explicit solution of the PY equation for a hard-sphere fluid in seven dimensions (7D). Such a solution allowed us to carry out an analysis of the virial coefficients arising both in the virial and in the compressibility routes and to determine the radius of convergence of both virial series. The results indicate some peculiar behavior of the virial coefficients with the virial route incorrectly predicting a negative fourth virial coefficient. The radius of convergence of the virial series is due to a singularity (branch point) located on the negative real axis and therefore what one has is an alternating series. Because of the good agreement between our value of the radius of convergence of the virial series and other independent estimates and the similar results obtained for $d=5$, it is tempting to conjecture that the PY solution for even higher dimensionalities should provide a rather accurate estimate of the radius of convergence of the true virial series and that it is the existence of singularities on the negative real axis (either poles or branch points) which determines such radius.

As a final point it is worth commenting that in this case our analysis was facilitated by the fact that we could combine both the analytical and the simulation results. And due to the common features such as the freezing transition that hard-core systems in different dimensionalities share, the expectation and the hope is that the present results shed some more light on the thermodynamic properties of such systems. As far as the high dimensionality limit is concerned, our results provide some support to the scenario of Frisch and Percus mentioned in the Introduction in the following sense. The solution to the PY equation predicts an alternating virial series. Further, the values of the scaled density $\hat{\rho}$ that one obtains for the radius of convergence ($\hat{\rho}=1.13$ for $d=5$, $\hat{\rho}=1.04$ for $d=7$, and the number $\hat{\rho}\approx 1.02$ coming out of our preliminary calculations for $d=9$) are consistent with a limiting value of $\hat{\rho}=1$ for $d\rightarrow\infty$. Also, the fluid range in $d=7$ is reasonably well accounted for by the first three or four virial coefficients so that it is conceivable that for infinite dimensionality only the second virial coefficient will be the dominant term.

ACKNOWLEDGMENTS

M.R. acknowledges the financial support of CONACyT through project No. I38644-E. M.L.H. acknowledges the hospitality of Universidad Complutense de Madrid (Spain), where the final version of the paper was prepared, as well as the financial support of DGAPA-UNAM during his sabbatical stay in Madrid. The research of A.S. has been partially

supported by the Ministerio de Ciencia y Tecnología (Spain) through grant No. BFM2001-0718 and by the European Community's Human Potential Program under contract HPRN-CT-2002-00307, DYGLAGEMEM.

APPENDIX A: SOLUTION OF THE PERCUS–YEVICK EQUATION FOR HARD HYPERSPHERES

For simplicity, in the remainder of this Appendix we set $\sigma=1$.

In the Percus–Yevick approximation, the structure factor $S(q)$ of a hard-sphere fluid in $d=2k+1$ dimensions is

$$S(q) = \frac{1}{\tilde{Q}(q)\tilde{Q}(-q)}, \quad (\text{A1})$$

where

$$\begin{aligned} \tilde{Q}(q) &= 1 - \lambda \int_0^1 dr e^{iqr} Q(r), \\ \lambda &\equiv (2\pi)^k \rho = 2^{2k}(2k+1)!! \eta, \end{aligned} \quad (\text{A2})$$

$Q(r)$ having the form

$$Q(r) = \begin{cases} \sum_{m=0}^k Q_m (r-1)^{m+k}, & 0 \leq r \leq 1 \\ 0, & r \geq 1 \end{cases}. \quad (\text{A3})$$

The $k+1$ coefficients $\{Q_m\}$ are functions of the density determined by the two linear equations

$$(-1)^k = -k! 2^k Q_k + \lambda \sum_{m=0}^k (-1)^m \frac{Q_m}{k+m+1}, \quad k \geq 0, \quad (\text{A4})$$

$$\begin{aligned} (-1)^k &= -(k-1)! 2^{k-1} Q_{k-1} \\ &+ \lambda \sum_{m=0}^k (-1)^m \frac{Q_m}{k+m+2}, \quad k \geq 1, \end{aligned} \quad (\text{A5})$$

plus the $k-1$ nonlinear equations

$$\begin{aligned} Q^{(2m+1)}(0) &= \frac{1}{2} \lambda (-1)^{m+1} [Q^{(m)}(0)]^2 \\ &- \lambda \sum_{\nu=0}^{m-1} (-1)^\nu Q^{(\nu)}(0) Q^{(2m-\nu)}(0), \\ 0 &\leq m \leq k-2. \end{aligned} \quad (\text{A6})$$

Here $Q^{(\nu)}(r)$ represents the ν th derivative of the function $Q(r)$. For $k=0$ ($d=1$), Eq. (A4) gives the exact solution for hard rods. For $k=1$ ($d=3$), Eqs. (A4) and (A5) are sufficient to find the solution of the PY equation. However, for $k \geq 2$ ($d \geq 5$) one needs in addition Eq. (A6), so that the problem reduces to solving an algebraic equation which, as we will argue below, is likely to be of degree $2^{k-1} = 2^{(d-3)/2}$.

In the limit $\eta \rightarrow 0$, it is easy to verify that

$$\lim_{\eta \rightarrow 0} Q_m = (-1)^{k+1} \frac{2^{-m}}{k!} \binom{k}{m}, \quad (\text{A7})$$

$$\lim_{\eta \rightarrow 0} Q(r) = (-1)^{k+1} \frac{2^{-k}}{k!} (r^2 - 1)^k,$$

$$\lim_{\eta \rightarrow 0} Q^{(2m)}(0) = (-1)^{m+1} 2^{-k} \frac{(2m)!}{m!(k-m)!}, \quad (\text{A8})$$

$$\lim_{\eta \rightarrow 0} Q^{(2m+1)}(0) = 0.$$

In general, one can expand the coefficients Q_m in powers of η :

$$Q_m(\eta) = \sum_{n=0}^{\infty} Q_{m,n} \eta^n, \quad (\text{A9})$$

where $Q_{m,0}$ is given by the first equation of (A7). Of course, the full nonlinear dependence of the coefficients $Q_m(\eta)$ can be obtained from the solution to the set of equations (A4)–(A6), either analytically ($k \leq 3$) or numerically ($k \geq 4$).

Once one has determined the function $Q(r)$, the structural properties of the fluid are given by Eqs. (A1) and (A2). In particular, the long wavelength limit of the structure factor and the contact value of the radial distribution function are, respectively,

$$S(q=0) = \frac{1}{[k! 2^k Q_k]^2}, \quad (\text{A10})$$

$$g(1^+) = (-1)^{k+1} k! Q_0. \quad (\text{A11})$$

The virial route to the EOS is given by

$$Z = 1 + 2^{d-1} \eta g(1^+), \quad (\text{A12})$$

while the compressibility route is

$$\chi \equiv k_B T \left(\frac{\partial \rho}{\partial p} \right)_T = S(q=0). \quad (\text{A13})$$

Inserting the expansion (A9) into Eqs. (A12) and (A13) we get the virial coefficients along both routes:

$$b_{n+2}^{\text{PY-v}} = 2^{2k} (-1)^{k+1} k! Q_{0,n}, \quad (\text{A14})$$

$$b_{n+1}^{\text{PY-c}} = 2^{2k} (k!)^2 \frac{1}{n+1} \sum_{m=0}^n Q_{k,m} Q_{k,n-m}.$$

1. The case $d=7$

Now we particularize to the seven-dimensional case ($k=3$), the unknowns being Q_m , $m=0, 1, 2, 3$. Since the two nonlinear equations (A6) involve the derivatives $Q^{(m)} \equiv Q^{(m)}(0)$, it is more advantageous to work with the set $\{Q^{(m)}\}$ rather than with the set $\{Q_m\}$. The latter can be expressed in terms of the former as

TABLE III. Values of the coefficients $Q_{m,n}$, defined by Eq. (A9), for $n=0-6$.

n	$Q_{0,n}$	$Q_{1,n}$	$Q_{2,n}$	$Q_{3,n}$
0	0.166 666 667	0.250 000 000	0.125 000 000	0.020 833 333
1	3.010 416 667	7.888 020 833	5.812 500 000	1.333 333 333
2	-5.119 791 667	-59.313 802 083	-41.072 916 667	-6.541 666 667
3	5.395 897 352 $\times 10^2$	4.247 567 790 $\times 10^3$	3.201 932 292 $\times 10^3$	6.540 590 278 $\times 10^2$
4	-2.286 456 505 $\times 10^4$	-2.488 562 475 $\times 10^5$	-1.884 527 380 $\times 10^5$	-3.841 568 446 $\times 10^4$
5	1.172 728 501 $\times 10^6$	1.635 350 292 $\times 10^7$	1.241 794 603 $\times 10^7$	2.538 798 289 $\times 10^6$
6	-6.600 274 174 $\times 10^7$	-1.143 524 052 $\times 10^9$	-8.688 923 174 $\times 10^8$	-1.778 764 760 $\times 10^8$

$$\begin{pmatrix} Q_0 \\ Q_1 \\ Q_2 \\ Q_3 \end{pmatrix} = - \begin{pmatrix} 20 & 10 & 2 & \frac{1}{6} \\ 45 & 25 & \frac{11}{2} & \frac{1}{2} \\ 36 & 21 & 5 & \frac{1}{2} \\ 10 & 6 & \frac{3}{2} & \frac{1}{6} \end{pmatrix} \cdot \begin{pmatrix} Q^{(0)} \\ Q^{(1)} \\ Q^{(2)} \\ Q^{(3)} \end{pmatrix}. \quad (A15)$$

Equations (A4) and (A5), plus Eq. (A6) with $m=0$ yield

$$Q^{(1)} = -3360\eta Q^{(0)2}, \quad (A16)$$

$$Q^{(2)} = - \frac{1 + 96Q^{(0)}\{1 - 5\eta[3 + 112Q^{(0)}(3 - 10\eta)]\}}{8(1 - \eta)}, \quad (A17)$$

$$Q^{(3)} = \frac{8 - 15\eta + 192Q^{(0)}\{2 - \eta[53 + 280Q^{(0)}(3 - 10\eta)^2 - 100\eta]\}}{8(1 - \eta)^2}. \quad (A18)$$

Thus, the parameters $Q^{(1)}$, $Q^{(2)}$, and $Q^{(3)}$ are given as explicit quadratic functions of $Q^{(0)}=Q(0)$. Finally, insertion of Eqs. (A16)–(A18) into Eq. (A6) with $m=1$ leads to the quartic equation

$$\begin{aligned} &8 - 15\eta + 192Q^{(0)}\{2 - \eta[88 - 135\eta + 1960Q^{(0)} \\ &\quad \times [3 - 4\eta[9 - 10\eta + 240Q^{(0)}(1 - \eta) \\ &\quad \times (3 - 10\eta(1 - 84Q^{(0)}(1 - \eta))]]]\} = 0. \end{aligned} \quad (A19)$$

Although an explicit expression exists for the physical root of Eq. (A19), it is of course too cumbersome and will be omitted here.³⁸

Table III shows the first few coefficients $Q_{m,n}$. The exact values are rational numbers, but they become more and more involved as the order n increases and so they are expressed in real form in Table III. From Eq. (A14) we can obtain the virial coefficients corresponding to the virial and the compressibility routes. The first few values are listed in Table I.

2. The case $d=9$

We will now sketch the result for the case $d=9$ following the same procedure. For $k=4$, the set $\{Q_m\}$ can be expressed in terms of the set $\{Q^{(m)}\}$ as

$$\begin{pmatrix} Q_0 \\ Q_1 \\ Q_2 \\ Q_3 \\ Q_4 \end{pmatrix} = \begin{pmatrix} 70 & 35 & \frac{15}{2} & \frac{5}{6} & \frac{1}{24} \\ 224 & 119 & 27 & \frac{19}{6} & \frac{1}{6} \\ 280 & 154 & \frac{73}{2} & \frac{9}{2} & \frac{1}{4} \\ 160 & 90 & 22 & \frac{17}{6} & \frac{1}{6} \\ 35 & 20 & 5 & \frac{2}{3} & \frac{1}{24} \end{pmatrix} \cdot \begin{pmatrix} Q^{(0)} \\ Q^{(1)} \\ Q^{(2)} \\ Q^{(3)} \\ Q^{(4)} \end{pmatrix}. \quad (A20)$$

In addition, the fifth derivative $Q^{(5)}$ is a linear combination of $\{Q_m\}$ and hence of the first four derivatives:

$$Q^{(5)} = -20(336Q^{(0)} + 210Q^{(1)} + 60Q^{(2)} + 10Q^{(3)} + Q^{(4)}). \quad (A21)$$

The nonlinear equations (A6) with $m=0, 1, 2$ allow one to express the odd derivatives in terms of the even ones as

$$Q^{(1)} = -\frac{\lambda}{2} Q^{(0)2}, \quad (A22)$$

$$Q^{(3)} = \frac{\lambda^3}{8} Q^{(0)4} - \lambda Q^{(0)} Q^{(2)}, \quad (\text{A23})$$

$$Q^{(5)} = -\frac{\lambda^5}{16} Q^{(0)6} + \frac{\lambda^3}{2} Q^{(0)3} Q^{(2)} - \frac{\lambda}{2} Q^{(2)2} - \lambda Q^{(0)} Q^{(4)}, \quad (\text{A24})$$

where $\lambda = 241\,920\eta$. Next, insertion of Eqs. (A22) and (A23) into the linear equations (A4) and (A5) yields $Q^{(2)}$ and $Q^{(4)}$ as *nonlinear* functions of $Q^{(0)}$. Finally, by equating the right-hand sides of Eqs. (A21) and (A24) one gets a closed algebraic equation of eighth degree for $Q^{(0)}$. A preliminary analysis of this equation indicates that its physical solution possesses a branch point at $\eta_{\text{branch}} \simeq -0.002\,394\,5$, so that the radius of convergence of the PY virial series would be $\eta_{\text{conv}} = |\eta_{\text{branch}}| \simeq 0.002\,394\,5$.

We have checked that for $d=11$ the resulting equation is of degree 16. Therefore, it seems plausible that in the general case $d=2k+1$ the degree of the equation for $Q^{(0)}$ is $2^{k-1} = 2^{(d-3)/2}$.

¹C. Freasier and D. J. Isbister, *Mol. Phys.* **42**, 927 (1981).

²M. Luban and A. Baram, *J. Chem. Phys.* **76**, 3233 (1982).

³C. G. Joslin, *J. Chem. Phys.* **77**, 2701 (1982).

⁴E. Leutheusser, *Physica A* **127**, 667 (1984).

⁵J. P. J. Michels and N. J. Trappeniers, *Phys. Lett. A* **104**, 425 (1984).

⁶H. L. Frisch, N. Rivier, and D. Wyler, *Phys. Rev. Lett.* **54**, 2061 (1985); **56**, 2331 (1986); M. Luban, *ibid.* **56**, 2330 (1986).

⁷W. Klein and H. L. Frisch, *J. Chem. Phys.* **84**, 968 (1986).

⁸J. L. Colot and M. Baus, *Phys. Lett. A* **119**, 135 (1986).

⁹M. Baus and J. L. Colot, *Phys. Rev. A* **36**, 3912 (1987).

¹⁰Y. Song, E. A. Mason, and R. M. Strat, *J. Phys. Chem.* **93**, 6916 (1989).

¹¹J. Amorós, J. R. Solana, and E. Villar, *Phys. Chem. Liq.* **19**, 119 (1989).

¹²Y. Song and E. A. Mason, *J. Chem. Phys.* **93**, 686 (1990).

¹³M. Luban and J. P. J. Michels, *Phys. Rev. A* **41**, 6796 (1990).

¹⁴D. J. González, L. E. González, and M. Silbert, *Phys. Chem. Liq.* **22**, 95 (1990).

¹⁵M. J. Maeso, J. R. Solana, J. Amorós, and E. Villar, *Mater. Chem. Phys.* **30**, 39 (1991).

¹⁶D. J. González, L. E. González, and M. Silbert, *Mol. Phys.* **74**, 613 (1991).

¹⁷L. E. González, D. J. González, and M. Silbert, *J. Chem. Phys.* **97**, 5132 (1992).

¹⁸E. Velasco, L. Mederos, and G. Navascués, *Mol. Phys.* **97**, 1273 (1999).

¹⁹H. L. Frisch and J. K. Percus, *Phys. Rev. E* **60**, 2942 (1999).

²⁰M. Bishop, A. Masters, and J. H. R. Clarke, *J. Chem. Phys.* **110**, 11449 (1999).

²¹A. Santos, *J. Chem. Phys.* **112**, 10680 (2000).

²²S. B. Yuste, A. Santos, and M. López de Haro, *Europhys. Lett.* **52**, 158 (2000).

²³G. Parisi and F. Slanina, *Phys. Rev. E* **62**, 6554 (2000).

²⁴A. Santos, S. B. Yuste, and M. López de Haro, *Mol. Phys.* **99**, 1959 (2001).

²⁵M. González-Melchor, J. Alejandro, and M. López de Haro, *J. Chem. Phys.* **114**, 4905 (2001).

²⁶R. Finken, M. Schmidt, and H. Löwen, *Phys. Rev. E* **65**, 016108 (2002).

²⁷E. Enciso, N. G. Almarza, M. A. González, and F. J. Bermejo, *Mol. Phys.* **100**, 1941 (2002).

²⁸N. Clisby and B. M. McCoy, *J. Stat. Phys.* **114**, 1343 (2004).

²⁹N. Clisby and B. M. McCoy, *J. Stat. Phys.* **114**, 1361 (2004).

³⁰N. Clisby and B. M. McCoy, *New results for the virial coefficients of D-dimensional hard spheres*, cond-mat/0303101 (2003).

³¹J. P. Hansen and I. R. McDonald, *Theory of Simple Liquids* (Academic, London, 1986).

³²So far, however, all known virial coefficients for hard rods, hard disks, and hard spheres are positive. For explicit values in the two latter cases see F. H. Ree and W. G. Hoover, *J. Chem. Phys.* **40**, 939 (1964); **46**, 4181 (1967); K. W. Kratky, *Physica A* **85**, 607 (1976); **87**, 584 (1977); *J. Stat. Phys.* **27**, 533 (1982); **29**, 129 (1982); E. J. Janse van Rensburg, *J. Phys. A* **26**, 4805 (1993); E. J. Janse van Rensburg and G. M. Torrie, **26**, 943 (1993).

³³In the notation of Ref. 23, $\ln \hat{\rho}^2 \rightarrow \rho_1 = d^{-1} \ln \bar{\rho}^2$, $\kappa \rightarrow \hat{\rho}_c$, $\hat{\rho}^{2d} \Delta \rightarrow P_c = \exp(dP_1/2)$, and $P_1 = 2 - 2 \ln 2 + \ln \hat{\rho}_c^2$.

³⁴In their paper (see Ref. 23), Parisi and Slanina do not locate the breakdown of Eq. (3) at $\hat{\rho} = \hat{\rho}_0$, but at a smaller value $\hat{\rho} = \hat{\rho}'_0 = [e/2 + \sqrt{(e/2)^2 - 1}]^{1/2} \exp[-\sqrt{1 - (2/e)^2}/2] \simeq 1.07\,614$. This value, which corresponds to $\kappa = 2/e$, is the density beyond which $\hat{\rho}^{2d} \Delta(\hat{\rho})$ diverges. The quantity $\hat{\rho}^{2d} \Delta(\hat{\rho})$ is the *absolute* correction to the *pressure*: $p \propto \hat{\rho}^d + \frac{1}{2} \hat{\rho}^{2d} + \hat{\rho}^{2d} \Delta(\hat{\rho})$. However, since the second virial term $\hat{\rho}^{2d}$ diverges as well in that limit, the relevant behavior is that of the *relative* correction $\Delta(\hat{\rho})$, which diverges at $\hat{\rho} = \hat{\rho}_0$.

³⁵N. F. Carnahan and K. E. Starling, *J. Chem. Phys.* **51**, 635 (1969).

³⁶As mentioned in Section I, a similar situation takes place with the PY solution for $d=5$, where $B_6^{\text{PY-v}} < 0$ but $B_6^{\text{PY-c}} > 0$.

³⁷M. P. Allen and D. J. Tildesley, *Computer Simulation of Liquids* (Clarendon, Oxford, 1987).

³⁸A code written in MATHEMATICA which provides the explicit expression for this physical root is available from any of the authors upon request.