



Erratum

Erratum to: “The penetrable-sphere fluid in the high-temperature, high-density limit” [Phys. Lett. A 323 (2004) 427]

Luis Acedo^a, Andrés Santos^{b,*}^a Instituto Universitario de Matemática Multidisciplinar, Universitat Politècnica de València, 46022 Valencia, Spain^b Departamento de Física, Universidad de Extremadura, 06071 Badajoz, Spain

ARTICLE INFO

ABSTRACT

Article history:

Received 20 April 2012

Accepted 22 May 2012

Available online 26 May 2012

Communicated by A.R. Bishop

A few misprints appearing in the paper [L. Acedo, A. Santos, Phys. Lett. A 323 (2004) 427] are corrected.

© 2012 Elsevier B.V. All rights reserved.

Keywords:

Penetrable-sphere model

Soft interactions

Spinodal instability

Fluid-solid phase transition

We have recently noticed a few misprints in Ref. [1], which, on the other hand, do not affect the physical results. The misprints are:

- Eq. (14) should be replaced by

$$\frac{s_{\text{ex}}^{\text{solid}}}{k_B} = d \ln \left[1 - \left(\frac{\hat{\eta}}{\alpha \eta_{\text{cp}}} \right)^{1/d} \right] + d \ln 2.$$

- Two lines below Eq. (14) of Ref. [1], $a_{\text{ex}}^{\text{solid}} = u_{\text{ex}}^{\text{solid}} + T s_{\text{ex}}^{\text{solid}}$ should be replaced by $a_{\text{ex}}^{\text{solid}} = u_{\text{ex}}^{\text{solid}} - T s_{\text{ex}}^{\text{solid}}$.
- Twelve lines below Fig. 3 of Ref. [1], $\mu_{\text{ex}}^{\text{fluid}}(\hat{\eta}_f) = \mu_{\text{ex}}^{\text{solid}}(\hat{\eta}_m)$ should be replaced by $\mu_{\text{ex}}^{\text{fluid}}(\hat{\eta}_f) = \mu_{\text{ex}}^{\text{solid}}(\hat{\eta}_m)$.
- The sixth and eighth columns of Table 2 of Ref. [1] need slight changes. The complete correct table is displayed as Table 1 of this Erratum.
- The shaded region in Fig. 3 of Ref. [1] should be slightly narrower. The correct figure is displayed as Fig. 1 of this Erratum.
- In Ref. [20b] of Ref. [1], 0161008 should be replaced by 016108.

Table 1

Values of the HS close-packing fraction η_{cp} , the wavenumber k_0 , the nearest-neighbor distance r_0 , the (scaled) spinodal instability packing fraction $\hat{\eta}_0$, the (scaled) freezing packing fraction $\hat{\eta}_f$, the (scaled) packing fraction $\hat{\eta}_{\text{ms}}$ at the condition of marginal stability and the (scaled) melting packing fraction $\hat{\eta}_m$.

d	η_{cp}	$k_0 \sigma$	r_0 / σ	$\hat{\eta}_0$	$\hat{\eta}_f$	$\hat{\eta}_{\text{ms}}$	$\hat{\eta}_m$
1	1	4.49	1.40	2.30	1.00	1.00	1.00
2	$\sqrt{3}\pi/6 \simeq 0.91$	5.14	1.37	1.89	0.90	0.95	1.01
3	$\sqrt{2}\pi/6 \simeq 0.74$	5.76	1.34	1.45	0.63	0.69	0.77
4	$\pi^2/16 \simeq 0.62$	6.38	1.32	1.07	0.37	0.41	0.48
5	$\sqrt{2}\pi^2/30 \simeq 0.47$	6.99	1.30	0.76	0.22	0.26	0.32

DOI of original article: <http://dx.doi.org/10.1016/j.physleta.2004.02.039>.

* Corresponding author.

E-mail addresses: luiacrod@imm.upv.es (L. Acedo), andres@unex.es (A. Santos).URL: <http://www.unex.es/fisteor/andres/> (A. Santos).




Fig. 1. Excess free energy per particle in the three-dimensional PS solid, $a_{\text{ex}}^{\text{solid}}/k_B T$ (solid line), and PS fluid, $a_{\text{ex}}^{\text{fluid}}/k_B T$ (dashed line), in the high-temperature limit. The excess internal energy, $u_{\text{ex}}^{\text{solid}}/k_B T$ (dotted line), and the excess entropy, $s_{\text{ex}}^{\text{solid}}/k_B$ (dash-dotted line), of the PS solid are also plotted. The shaded area represents the fluid–solid coexistence region.

References

- [1] L. Acedo, A. Santos, Phys. Lett. A 323 (2004) 427.