

Does the Chapman–Enskog Expansion for Sheared Granular Gases Converge?

Andrés Santos*

Departamento de Física, Universidad de Extremadura, E-06071 Badajoz, Spain

(Received 21 November 2007; published 21 February 2008)

The fundamental question addressed in this Letter is whether or not the partial Chapman–Enskog expansion $P_{xy} = -\sum_{k=0}^{\infty} \eta_k (\partial u_x / \partial y)^{2k+1}$ of the shear stress converges for a gas of inelastic hard spheres. By using a simple kinetic model it is shown that, in contrast to the elastic case, the above series does converge, the radius of convergence increasing with inelasticity. It is argued that this paradoxical conclusion is not an artifact of the kinetic model and can be understood in terms of the time evolution of the scaled shear rate in the uniform shear flow.

DOI: [10.1103/PhysRevLett.100.078003](https://doi.org/10.1103/PhysRevLett.100.078003)

PACS numbers: 45.70.Mg, 05.20.Dd, 47.50.-d, 51.10.+y

The hydrodynamic description of conventional fluids is usually restricted to the Navier–Stokes (NS) constitutive equations [1]. For instance, if the flow is incompressible (i.e., $\nabla \cdot \mathbf{u} = 0$, where \mathbf{u} is the flow velocity), Newton’s law establishes a linear relationship $P_{xy} = -\eta_0 \partial u_x / \partial y$ between the shear stress P_{xy} and the shear rate $\partial u_x / \partial y$, where η_0 is the shear viscosity and it has been assumed that $\partial u_y / \partial x = 0$. The NS constitutive equations represent excellent approximations in most of the physical situations of experimental interest, even if the regime is turbulent [2]. In the case of a dilute gas, they can be justified under the assumption that the smallest of the characteristic hydrodynamic lengths (L) associated with the gradients of density (n), temperature (T), and flow velocity (\mathbf{u}) are much larger than the mean free path ℓ of the gas particles, i.e., $\mu \equiv \ell/L \ll 1$. As a matter of fact, the Chapman–Enskog (CE) method provides a systematic scheme to obtain the *normal* solution of the Boltzmann equation as an expansion in powers of the uniformity parameter (or Knudsen number) μ [3]. The leading terms in the CE expansion yield the NS constitutive equations, with the bonus of providing expressions for the transport coefficients (like the shear viscosity η_0) in terms of the microscopic properties and of the hydrodynamic quantities.

A fundamental question concerning the CE method is the nature (convergent versus divergent) of the expansion in powers of μ . More than 40 years ago Grad [4] provided compelling arguments on the asymptotic character of the CE expansion. This means that, as expected on physical grounds, the CE series truncated at a given order (e.g., NS, Burnett, super-Burnett, etc.) becomes closer and closer to the true value as μ becomes smaller and smaller. Of course, the asymptotic character of the CE expansion does not imply (but is implied by) the stronger condition of convergence. However, McLennan [5] was able to prove convergence of the *partial* sum of the CE series made of *linear* terms (i.e., terms of the form $\nabla_{i_1} \nabla_{i_2} \dots \nabla_{i_k} A$, where $A = n, T$, or \mathbf{u}) for a general class of cutoff potentials.

What about the *nonlinear* terms of the series? To be more specific, let us consider the following subclass of the

full CE series of the shear stress:

$$P_{xy} = -\sum_{k=0}^{\infty} \eta_k (\partial u_x / \partial y)^{2k+1}, \quad (1)$$

where η_0 is the NS shear viscosity, η_1 is a super-Burnett coefficient, and so on. The full CE series of P_{xy} reduces to the partial series (1) if (and only if) $\partial u_i / \partial x_j = \dot{\gamma} \delta_{ix} \delta_{jy}$ and $\nabla n = \nabla T = \nabla \dot{\gamma} = \mathbf{0}$; i.e., the only nonzero hydrodynamic gradient is a uniform shear rate $\dot{\gamma} \equiv \partial u_x / \partial y$. Interestingly enough, there exists a physical state (the so-called simple or uniform shear flow, USF) that is consistent with those conditions [6,7]. In that state, the density and the shear rate are constant in time, but the temperature increases due to viscous heating. The identification of the characteristic hydrodynamic length L is (except for a numerical factor) unambiguous: $L \sim \sqrt{2T/m}/\dot{\gamma}$ (where m is the mass of a particle), so that the uniformity parameter becomes $\mu = \dot{\gamma}/\nu$, where $\nu \sim \sqrt{2T/m}/\ell$ is a characteristic collision frequency. Thus, the CE expansion (1) can be recast into the dimensionless form

$$P_{xy}/nT = -\mu F(\mu^2), \quad F(z) = \sum_{k=0}^{\infty} c_k z^k, \quad (2)$$

where $c_k \equiv (\eta_k/nT)\nu^{2k+1}$. Despite its simple definition, the USF state is complex enough to prevent an exact solution of the nonlinear Boltzmann equation. However, the problem becomes solvable in the framework of the Bhatnagar–Gross–Krook (BGK) model kinetic equation [7], and the solution shows that, for a wide class of repulsive potentials, the CE expansion (1) is *divergent* [8]. More specifically, the dimensionless coefficients c_k behave for large k as $|c_k| \sim (2/d)^k k!$ in the case of d -dimensional hard spheres.

In the preceding paragraphs we have considered conventional gases made of particles that collide elastically. On the other hand, the same issues discussed above can be applied to granular gases, i.e., large assemblies of (mesoscopic or macroscopic) particles that collide inelastically and are maintained in fluidized states. Apart from their

practical interest, granular gases are physical systems worth studying at a fundamental level because they are intrinsically out of equilibrium and exhibit a wide repertoire of complex and exotic behavior [9,10]. Nonequilibrium statistical-mechanical concepts and tools, in particular, the kinetic theory approach based on the Boltzmann and Enskog equations suitably modified to account for inelastic collisions, have proven to be useful to understand the behavior of granular gases [11]. Most of the theoretical efforts have focused on a minimal model of granular gases consisting of smooth inelastic hard spheres characterized by a constant coefficient of normal restitution $\alpha < 1$. Specifically, the CE method has been applied to the (inelastic) Boltzmann and Enskog kinetic equations and the NS transport coefficients have been derived [12].

The question I want to address in this Letter is, does the nonlinear subclass of the full CE expansion converge in the case of a gas of inelastic hard spheres? Given that the answer is negative when the gas is made of elastic hard spheres ($\alpha = 1$) [8], the strong challenges to the validity of hydrodynamics in granular media [10], and the inherently non-Newtonian nature of the steady USF of granular gases [13], it seems plausible to expect that the (partial) CE series (1) is divergent for granular gases. It will be shown below that, paradoxically, the series (1) does converge in the case of inelastic hard spheres and that, in fact, the radius of convergence increases with increasing inelasticity.

The energy balance equation for inelastic hard spheres in the USF is

$$\partial_t T(t) = -(2/dn)\dot{\gamma}P_{xy}(t) - \zeta(t)T(t), \quad (3)$$

where ζ is the so-called cooling rate [11]. From the Boltzmann equation it is possible to show that it is approximately given by $\zeta = \frac{d+2}{4d}(1-\alpha^2)\nu_0$, ν_0 being an effective collision frequency for elastic spheres. The cooling term on the right-hand side of Eq. (3) competes with the viscous heating term, so that, depending on the initial state and the value of $\dot{\gamma}$, the temperature either grows or decreases with time until a steady state is eventually reached [13,14]. In order to relate the shear stress $P_{xy}(t)$ to the shear rate $\dot{\gamma}$ and to the temperature $T(t)$, and thus analyze the series representation (2), let us replace the Boltzmann equation by the BGK-like kinetic model [15]

$$(\partial_t + \mathbf{v} \cdot \nabla)f = -\nu(f - f_0) + (\zeta/2)\partial_v \cdot (\mathbf{V}f), \quad (4)$$

where f is the velocity distribution function, $\mathbf{V} \equiv \mathbf{v} - \mathbf{u}$ is the peculiar velocity, f_0 is the local version of the homogeneous cooling state distribution [11] (parametrized by the actual fields n , \mathbf{u} , and T), and ν is an effective collision frequency. Here the simple choice $\nu = \frac{1+\alpha}{2}\nu_0$ is made, so that $\epsilon \equiv \zeta/\nu = \frac{d+2}{2d}(1-\alpha)$. Taking moments in Eq. (4) one gets

$$\begin{aligned} \partial_t P_{xy}(t) &= -\dot{\gamma}P_{xy}(t) - [\nu(t) + \zeta(t)]P_{xy}(t), \\ \partial_t P_{yy}(t) &= n\nu(t)T(t) - [\nu(t) + \zeta(t)]P_{yy}(t), \end{aligned} \quad (5)$$

where P_{yy} is a normal stress. Note that the explicit form of f_0 is not needed in the derivation of Eq. (5) and so no assumption of f or f_0 being close to a Maxwellian is taken. Equation (5), with $\nu = \frac{1+\alpha}{2}[1 - \frac{d-1}{2d}(1-\alpha)]\nu_0$, is also obtained from the Boltzmann equation in Grad's moment approximation [13]. Equations (3) and (5) constitute a closed set of equations for the evolution of T , P_{xy} , and P_{yy} . In order to describe the hydrodynamic regime and analyze the CE expansion, we must focus on the nonlinear dependence of the scaled viscosity function $F(\mu^2(t)) \equiv -P_{xy}(t)/nT(t)\mu(t)$ as a function of the scaled shear rate (or uniformity parameter) $\mu(t) \equiv \dot{\gamma}/\nu(t)$. Elimination of time in favor of $z \equiv \mu^2 \propto 1/T$ in Eqs. (3) and (5) yields the following single second-order ordinary differential equation:

$$\begin{aligned} 1 - \left(1 + \frac{2}{d}zF\right)^2 F &= \frac{1}{2}\left(\epsilon - \frac{2}{d}zF\right) \\ &\times \left\{z\partial_z \left[\left(\epsilon - \frac{2}{d}zF\right)(F + 2z\partial_z F) \right] \right. \\ &+ \left(1 + \frac{2}{d}zF\right)(F + 2z\partial_z F) \\ &\left. + 2z\partial_z \left[\left(1 + \frac{2}{d}zF\right)F \right] \right\}. \end{aligned} \quad (6)$$

Insertion of the series expansion $F(z) = \sum_{k=0}^{\infty} c_k z^k$ into Eq. (6) allows one to get a recursive relation expressing the coefficient c_k for $k \geq 1$ in terms of the lower-order coefficients $\{c_{k'}, 0 \leq k' \leq k-1\}$. With the help of a computer algebra system, one can obtain the coefficients c_k , for given d and α , as exact rational numbers up to a maximum value of k limited by computer time and internal memory. The results of c_k presented here have been obtained for $k \leq 400$.

The first few coefficients c_k are listed in Table I for $d = 3$ and $\alpha = 0.5, 0.7, 0.9, 0.99$, and 1 (elastic hard spheres). As is already apparent from Table I, the magnitude of the coefficients increases as the system becomes less inelastic, this effect being more and more dramatic with increasing k . For instance, the value of $|c_{20}|$ is 1.2×10^{-5} , 5.2×10^{-3} , 2.1×10^5 , 1.4×10^{11} , and 1.1×10^{12} for $\alpha = 0.5, 0.7, 0.9, 0.99$, and 1, respectively. Figure 1 shows a log-normal plot of $|c_k|$ for the four inelastic systems considered. The results are consistent with an asymptotic behavior of the form $|c_k| \approx Ak^{-a}z_s^{-k}$, where $z_s = \frac{d}{2}\epsilon(1+\epsilon)^2$ is the steady-state value of the square uniformity parameter. The exponent a and the amplitude A are estimated by a linear fit of a plot of $\ln(|c_k|z_s^k)$ versus $\ln k$ for $100 \leq k \leq 400$, the fit being less robust in the case $\alpha = 0.5$ than in the other cases. The numerical values of z_s , as well as of the estimates of a and $\ln A$, are also included in Table I. In principle, the viscosity function $F(z)$ is different for each dimensionality. However, inspection of Eq. (6) shows that, for a given value of ϵ , the dependence of F on z and d occurs only

TABLE I. CE coefficients c_k for $0 \leq k \leq 6$, steady-state value of the square uniformity parameter, z_s , and estimates of the parameters a and $\ln A$ associated with the asymptotic behavior of $|c_k|$.

α	c_0	c_1	c_2	c_3	c_4	c_5	c_6	z_s	a	$\ln A$
0.5	0.827 586	-0.380 155	0.067 708 5	0.040 419 6	-0.002 517 05	-0.011 976 7	-0.003 995 93	1.254 34	15.8	60.3
0.7	0.888 889	-0.536 326	0.146 688	0.110 743	-0.018 097 0	-0.072 768 1	-0.033 608 2	0.585 938	8.21	7.08
0.9	0.960 000	-0.798 195	0.377 471	0.376 385	-0.188 551	-0.778 076	-0.618 849	0.146 701	15.8	22.6
0.99	0.995 851	-0.976 777	0.627 438	0.720 350	-0.694 311	-3.200 37	-3.756 20	0.012 709 2	95.9	283
1	1	-1	0.666 667	0.777 778	-0.814 815	-3.827 16	-4.724 28	0

through the combination z/d . Therefore, seen as a function of ϵ and z/z_s , $F(z)$ is independent of d .

The large- k asymptotic behavior $\ln|c_k| \sim -k \ln z_s$ for $\alpha < 1$ implies that the CE series (1) is *convergent* for inelastic hard spheres. Moreover, the radius of convergence is $\partial u_x / \partial y = \nu z_s^{1/2}$ and thus increases with inelasticity. Equation (6) shows that $z = z_s$ is indeed a singular point since the steady state (z_s, F_s) , with $F_s = (1 + \epsilon)^{-2}$, is the solution of $1 - (1 + \frac{2}{3} z F)^2 F = \epsilon - \frac{2}{3} z F = 0$. What is relevant here is that $z = 0$ is a regular point (except in the elastic case) and $z = z_s$ is the singularity of $F(z)$ in the complex z plane closest to the origin. The convergent character of the series (2) for the most inelastic case considered here ($\alpha = 0.5$) is illustrated in Fig. 2, where the numerical solution [13] of Eq. (6) is compared with the *truncated* CE series $F_N(z) \equiv \sum_{k=0}^N c_k z^k$ for several values of the truncation order N . One can observe that the truncated series agree among themselves and with the numerical solution for $z < z_s$. For $z > z_s$, however, the CE expansion becomes useless and one must determine $F(z)$ numerically from Eq. (6). An alternative method, successfully used in the elastic case [8], would consist of expanding $F(z)$ around the point at infinity as $F(z) = z^{-2/3} \sum_{k=0}^{\infty} \bar{c}_k z^{-k/3}$. Figure 2 also includes results recently obtained from Monte Carlo simulations of the Boltzmann equation [14], which show that the predictions of Eq. (6) are quantitatively accurate, even for strong inelasticity, except that Eq. (6) underestimates the location of the steady-state point (z_s, F_s) . The simulation curve shown in Fig. 2 is the collapse of data obtained by the direct simu-

lation Monte Carlo method starting from 20 different initial conditions (ten for $z < z_s$ and ten for $z > z_s$), letting the system evolve in time, and discarding the kinetic transients lasting a few mean free times. This explains why the simulation curve does not reach the point $z = 0$. For simulation details the reader is referred to Ref. [14].

Is the paradoxical regularization by inelasticity of the CE series (1) an artifact of the kinetic model? The following physical argument suggests that this is not the case. Let us assume that the reference homogeneous state is perturbed at time $t = 0$ by a *weak* shear rate $\dot{\gamma}$. Thus the homogeneous state ($\mu = 0$) is stable or unstable against this USF perturbation depending on whether the time-dependent uniformity parameter $\mu(t) = \dot{\gamma}/\nu(t)$ asymptotically goes to zero or grows with time. The first situation occurs in a conventional gas of elastic hard spheres since the viscous heating produces a monotonic increase of temperature [cf. Eq. (3) with $\zeta = 0$] and thus $\mu(t) \rightarrow 0$. However, in a gas of inelastic hard spheres it is always possible that the perturbation is small enough to make the cooling rate prevail over the viscous heating rate and so the temperature keeps decreasing in time [cf. Eq. (3) with $\zeta > 0$] until the steady state is eventually reached; thus the




FIG. 1 (color online). Plot of $\text{sgn}(c_k) \ln|c_k|$ for $d = 3$ and $\alpha = 0.5, 0.7, 0.9$, and 0.99 . The thin solid lines represent $\pm(-k \ln z_s - a \ln k + \ln A)$, where a and $\ln A$ are fitting parameters.




FIG. 2 (color online). Scaled viscosity function $F(z)$ for $d = 3$ and $\alpha = 0.5$. The thin solid line represents the numerical solution of Eq. (6), while the dashed lines represent the truncated CE expansion $F_N(z)$ with $N = 5, 20, 50, 100$, and 200 . The thick solid line represents the results obtained from Monte Carlo simulations of the Boltzmann equation [14]. The steady-state points (z_s, F_s) obtained from the kinetic model and the simulations are indicated by the triangle and the circle, respectively. The inset includes $F_N(z)$ with $N = 1-4$ in the region $0 \leq z \leq 1.5$.

uniformity parameter grows, i.e., $\mu(t) \rightarrow z_s^{1/2}$. Since the CE expansion (2) measures the departure from the reference homogeneous state ($\mu = 0$), it is reasonable to expect that the series diverges if $\mu(t)$ goes to zero while it converges if $\mu(t)$ increases with time.

The above heuristic argument can also be applied to the simplest CE series in a compressible flow, namely,

$$P_{yy} - nT = - \sum_{k=0}^{\infty} \eta'_k (\partial u_y / \partial y)^{k+1}. \quad (7)$$

The full CE expansion of P_{yy} reduces to the series (7) in the uniform longitudinal flow (ULF) characterized by [16] $\partial u_y / \partial y = \dot{\gamma}(t) = \dot{\gamma}_0 / (1 + \dot{\gamma}_0 t)$ and $n(t) = (n_0 / \dot{\gamma}_0) \dot{\gamma}(t)$. The exact energy balance equation, $\partial_t T(t) = -(2/dn) \dot{\gamma}(t) P_{yy}(t) - \zeta(t) T(t)$, shows that the uniformity parameter $\mu(t) = \dot{\gamma}(t) / \nu(t)$ increases with time if $\dot{\gamma}_0 > 0$, both for elastic and inelastic collisions. However, if $\dot{\gamma}_0 < 0$, $|\mu(t)| \rightarrow 0$ in the elastic case, whereas $|\mu(t)|$ increases toward a stationary value in the inelastic case. Thus, the CE expansion (7) is expected to diverge for conventional gases and converge for granular gases. This expectation is confirmed by an analysis of the ULF based again on the kinetic model (4).

It is important to note that the dependence on n , T , and α of the transport coefficients η_k and η'_k in the partial CE expansions (1) and (7), respectively, is not influenced by the specific state under consideration. Accordingly, the series (1) and (7) converge or diverge irrespective of whether the system is in the USF, the ULF, or in any other state. The advantage of the USF and ULF is that the full CE series of the shear stress and the normal stress reduce to the partial series (1) and (7), respectively, thus allowing us to explore their character in a rather detailed way. Regarding the CE subseries made of linear terms, it is reasonable to expect that, as in the elastic case, it also converges for inelastic collisions, especially if one takes into account the exact mapping between the inelastic and elastic versions of the Enskog–Lorentz equations [17].

The CE expansion is the main route to hydrodynamics, and so its convergence or divergence has a renewed interest in granular gases in view of some debate on the applicability of hydrodynamics to this class of nonequilibrium systems [10]. I expect that this Letter can contribute to a clarification of this controversial issue by presenting a case study where the application of the CE expansion to describe the nonlinear regime might have a larger practical interest in granular than in conventional gases.

I am grateful to J.W. Dufty and V. Garzó for useful comments on an early version of this Letter. This work has been supported by the Ministerio de Educación y Ciencia (Spain) through Grant No. FIS2007–60977 (partially financed by FEDER funds) and by the Junta de Extremadura

(Spain) through Grant No. GRU07046.

*Electronic addresses: andres@unex.es
http://www.unex.es/fisteor/andres/

- [1] S.R. de Groot and P. Mazur, *Nonequilibrium Thermodynamics* (Dover, New York, 1984).
- [2] R.J. Donnelly, in *A Physicist's Desk Reference*, edited by H.L. Anderson (American Institute of Physics, New York, 1989), pp. 196–209.
- [3] S. Chapman and T.G. Cowling, *The Mathematical Theory of Non-uniform Gases* (Cambridge University Press, Cambridge, 1970).
- [4] H. Grad, *Phys. Fluids* **6**, 147 (1963).
- [5] J.A. McLennan, *Phys. Fluids* **8**, 1580 (1965); G. Scharf, *Helv. Phys. Acta* **40**, 929 (1967); **42**, 5 (1969).
- [6] A.W. Lees and S.F. Edwards, *J. Phys. C* **5**, 1921 (1972).
- [7] V. Garzó and A. Santos, *Kinetic Theory of Gases in Shear Flows. Nonlinear Transport* (Kluwer, Dordrecht, 2003), and references therein.
- [8] A. Santos, J.J. Brey, and J.W. Dufty, *Phys. Rev. Lett.* **56**, 1571 (1986); A. Santos and J.J. Brey, *Physica (Amsterdam)* **174A**, 355 (1991).
- [9] H.M. Jaeger, S.R. Nagel, and R. Behringer, *Phys. Today* **49**, No. 4, 32 (1996); *Rev. Mod. Phys.* **68**, 1259 (1996); J.M. Ottino and D.V. Khakhar, *Annu. Rev. Fluid Mech.* **32**, 55 (2000); A. Kudrolli, *Rep. Prog. Phys.* **67**, 209 (2004).
- [10] L. Kadanoff, *Rev. Mod. Phys.* **71**, 435 (1999).
- [11] N. Brilliantov and T. Pöschel, *Kinetic Theory of Granular Gases* (Oxford University Press, Oxford, 2004); C.S. Campbell, *Annu. Rev. Fluid Mech.* **22**, 57 (1990); J.W. Dufty, *J. Phys. Condens. Matter* **12**, A47 (2000); arXiv:0709.0479; I. Goldhirsch, *Annu. Rev. Fluid Mech.* **35**, 267 (2003).
- [12] J.J. Brey, J.W. Dufty, C.S. Kim, and A. Santos, *Phys. Rev. E* **58**, 4638 (1998); V. Garzó and J.W. Dufty, *Phys. Rev. E* **59**, 5895 (1999); *Phys. Fluids* **14**, 1476 (2002); V. Garzó, J.W. Dufty, and C.M. Hrenya, *Phys. Rev. E* **76**, 031303 (2007); V. Garzó, C.M. Hrenya, and J.W. Dufty, *Phys. Rev. E* **76**, 031304 (2007); S.H. Noskovicz, O. Bar-Lev, D. Serero, and I. Goldhirsch, *Europhys. Lett.* **79**, 60001 (2007).
- [13] A. Santos, V. Garzó, and J.W. Dufty, *Phys. Rev. E* **69**, 061303 (2004).
- [14] A. Astillero and A. Santos, *Europhys. Lett.* **78**, 24002 (2007).
- [15] J.J. Brey, J.W. Dufty, and A. Santos, *J. Stat. Phys.* **97**, 281 (1999).
- [16] A.N. Gorban and I.V. Karlin, *Phys. Rev. Lett.* **77**, 282 (1996); I.V. Karlin, G. Dukek, and T.F. Nonnenmacher, *Phys. Rev. E* **55**, 1573 (1997); A. Santos, *Phys. Rev. E* **62**, 6597 (2000).
- [17] A. Santos and J.W. Dufty, *Phys. Rev. Lett.* **97**, 058001 (2006).